

Box 217, Oakland City, Indiana 47660, USA.

TRACHEMYS VENUSTA (Mesoamerican Slider). USA: FLORIDA: GILCHRIST Co.: Santa Fe River, 1.2 km downstream from Rum Island (29.834354°N, 82.690575°W; datum WGS84). 19 January 2010. Matthew H. Kail. Verified by Kurt Buhmann and Michael Seidel. Florida Museum of Natural History (UF 157304). New state record. Adult male (straight carapace length 243 mm, plastron length 216 mm, mass 1690 g) captured by hand at 2130 h along the northern shoreline. High leech load (80–100 leeches) and presence of algae on carapace suggest that this is not a recently released captive. This non-native species may potentially harm the closely related native Yellow-bellied Slider (*Trachemys scripta scripta*) population through interbreeding and genetic introgression.

Submitted by **MATTHEW H. KAIL** (e-mail: snappergi@aol.com), and **DARCY T. GARRENTON**, Teaching Zoo, Santa Fe College, Gainesville, Florida 32606, USA (e-mail: dgarrent@kent.edu); **ERIC SUAREZ**, Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, Florida 32611, USA (e-mail: eric.suarez725@yahoo.com); **GERALD R. JOHNSTON**, Department of Natural Sciences, Santa Fe College, Gainesville, Florida 32606, USA (e-mail: jerry.johnston@sfcollge.edu); and **JOSEPH C. MITCHELL**, Mitchell Ecological Research Services, Gainesville, Florida 32627, USA (e-mail: dr.joe.mitchell@gmail.com).

SQUAMATA — LIZARDS

ANOLIS (=NOROPS) SAGREI (Cuban Brown Anole). USA: TEXAS: FORT BEND Co.: Collected in author's backyard (29.702888°N, 95.790955°W; WGS84). 10 March 2010. A. Wood. Texas Cooperative Wildlife Collection (TCWC 94268). Verified by James R. Dixon. New county record (Dixon 2000. *Amphibians and Reptiles of Texas*. 2nd ed., Texas A&M Univ. Press, College Station, 421 pp.)

Submitted by **ADAM WOOD**, 24018 Seventh Heaven, Katy, Texas 77494, USA; e-mail: saleenadam@hotmail.com.

ANOTOSAURA VANZOLINIA. BRAZIL: RIO GRANDE DO NORTE: Municipality of Tenente Laurentino Cruz, Serra Nova (06.109°S, 36.719°W; datum WGS84, elev 704 m). 21 September 2009; 21 and 23 October 2009. L. Barros Ribeiro. Coleção Herpetológica do Departamento de Botânica, Ecologia e Zoologia, Universidade Federal do Rio Grande do Norte, Natal. Rio Grande do Norte (CHBEZ 2863, 2924, 2925). Verified by M. T. Rodrigues. First state record, extends the known geographical distribution of this species ca. 135 km N from the nearest record in the municipalities of São José dos Cordeiros/Sumé (Reserva Particular do Patrimônio Natural Fazenda Almas) (07.471°S, 36.881°W), state of Paraíba.

The gymnophthalmid lizard *Anotosaura vanzolinia* has been found in Rui Barbosa, Bahia State (Freitas and Silva 2007. *Guia Ilustrado: a Herpetofauna das Caatingas e Áreas de Altitudes do Nordeste Brasileiro*. Editora USEB, Pelotas, Brazil. 384 pp.); Agrestina (type locality) and district of Serra Negra, Bezerras municipality, state of Pernambuco (Rodrigues 1986. *Pap. Avul. Zool.* 36[20]: 237–250); Cabaceiras (Rodrigues 1986, *op. cit.*); São José da Mata (Delfim and Freire 2007. *Oecol. Bras.* 11[3]:365–382), São

José dos Cordeiros and Sumé (Reserva Particular do Patrimônio Natural Fazenda Almas) in the state of Paraíba (Freire et al. 2009. *In* E. M. X. Freire [org.], *Répteis Squamata das Caatingas do Seridó do Rio Grande do Norte e do Cariri da Paraíba: Síntese do Conhecimento Atual e Perspectivas*, pp. 51–84. Editora da UFRN, Natal, RN, Brazil).

Submitted by **MELISSA GOGLIATH** (e-mail: melbiologa@gmail.com)^{1,2}, **LEONARDO B. RIBEIRO** (e-mail: ribeiro.lb@gmail.com)^{1,2}, and **ELIZA M. X. FREIRE** (e-mail: elizajuju@ufrnet.br)^{1,2}, ¹Laboratório de Herpetologia, Departamento de Botânica, Ecologia e Zoologia, Centro de Biociências, Universidade Federal do Rio Grande do Norte, Campus Universitário, 59072-970, Natal, RN, Brazil; ²Programa de Pós-graduação em Psicobiologia/ Universidade Federal do Rio Grande do Norte, 59078-970, Natal, RN, Brazil.

ASPIDOSCELIS SEXLINEATA (Six-lined Racerunner). USA: ILLINOIS: HANCOCK Co.: One *Aspidoscelis sexlineata* was captured on the Mississippi River sand levee south of Warsaw, Illinois (40.34295°N, 91.44906°W; WGS 84). 09 August 2009. James T. Lamer and Terri L. Tobias. Illinois Natural History Survey (INHS) 21467. Verified by Chris Phillips. New county record (Phillips et al. 1999. *Field Guide to Amphibians and Reptiles of Illinois*. Illinois Nat. Hist. Surv. Manual 8, 282 pp.). Four other individuals were observed in the area.

Submitted by **TERRI L. TOBIAS** and **JAMES T. LAMER**, Western Illinois University, Alice Kibbe Field Station, Warsaw, Illinois 62379, USA.

CHAMAELEO GRACILIS (Slender Chamaeleon). MALI: SI-KASSO REGION: 8 km E of Yanfolila (11.11°N, 08.04°W; no datum available). 05 July 2008. Sébastien Trape. Institut de Recherche pour le Développement at Dakar (IRD TR-2953). Verified by Laurent Chirio. First record for Mali where only *Chamaeleo senegalensis* and *C. africanus* were previously known (Joger and Lambert 1996. *In* Ulrich [ed.], *Tropical Biodiversity and Systematics*. Proceeding of the International Symposium on Biodiversity and Systematics in Tropical Ecosystems, Bonn, 1994, pp. 189–202. Zoologisches Forschungsinstitut und Museum Alexander Koenig, Bonn; TIGR Reptile Database 2009. <http://www.reptile.database.org/>. Accessed December 2009).

Submitted by **SEBASTIEN TRAPE**, Laboratoire ECOLAG, Université Montpellier II, 34095 Montpellier Cedex 5, France; e-mail: sebastien_trape@yahoo.fr.

DAREVSKIA BRAUNERI SZCZERBAKI (Rock Lizard). GEORGIA: ABKHAZIA: GUDAUTA DISTRICT: Lzaa village, Pitsunda-Myussera Hills. ZISP 24397, 25816–25818. 17 April 2006. Three specimens found by Konstantin Milto and Mark Pestov. Several specimens were registered in 5–7 July, 7–9 August 2008, and 7–12 August 2009 by Mark Pestov and Olga Bezman-Moseyko. Verified by Ilya S. Darevsky. Lizards observed on clay-rock cliffs of Black Sea coast between Lzaa settlement (43.16813°N, 40.4108°E; no datum available) and mouth of Mysra River (43.15009°N, 40.45921°E). Known from a small territory on northeastern coast of Black Sea in Krasnodar Region, Russia, and a subspecies endemic to Crimea-Novorossisk subprovince of East Mediterranean Province in Caucasus, inhabiting narrow coastal zone from Anapa

town to Cape of Utrish. This new locality, 297 km to southeast from earlier localities, may be the southernmost border of the range, and is a first record for the country.

Submitted by **KONSTANTIN D. MILTO** (e-mail: coluber@zin.ru), **MARK V. PESTOV**, and **OLGAS. BEZMAN-MOSEYKO**, Zoological Institute, Department of Herpetology, St. Petersburg, Universitetskaya emb., 1, 199034, Russia.

MESALINA BREVIROSTRIS (Short-nosed Desert Racer). IRAN: SISTAN & BALUCHISTAN PROVINCE: Gando Protected Area: Garm Beet (61.30°N, 25.47°E; ca. 146 m elev.). Nastaran Heydari. 20 October 2008. Verified by Steven C. Anderson. Zoology Museum Golestan University, ZMGU 2429. First confirmed record for Sistan and Baluchistan Province. Associated with *Phoenixdactylifera*, *Prosopis cineraria*, and *Nannorhops ritchieana* in lowland sandy hills. 850 km E of nearest record (Qeshm Island and Mahor Birinji; Anderson 1999. The Lizards of Iran. Society for the Study of Amphibians and Reptiles, Ithaca, New York. 415 pp.; Haas and Werner 1969. Bull. Mus. Comp. Zool., Harvard Univ. 138[6]:327–405).

Submitted by **NASTARAN HEYDARI**, Department of Biodiversity, Khuzestan University of Science and Research, Ahwaz, Iran (e-mail: heydari.pr@gmail.com); **HAJI GHOLI KAMI**, Department of Biology, Faculty of Sciences, Golestan University, Gorgan, Golestan Province, Iran (e-mail: hgkami2000@yahoo.com); and **SOHEILA SHAFIEI**, Department of Biology, Faculty of Science, Shahid Bahonar University of Kerman, Iran (e-mail: shafiei_soheila@yahoo.com).

PANASPIS NIMBAENSIS (Mount Nimba Lidless Skink). SENEGAL: ZIGUINCHOR REGION: Djibonker (12.32°N, 16.21°W; no datum available). 01 March 2009. Sébastien Trape. Institut de Recherche pour le Développement at Dakar (IRD TR-2954). Verified by Laurent Chirio. First record for Senegal (Cisse and Karns 1978. Bull. IFAN. 40A:144–211; TIGR Reptile Database 2009. <http://www.reptile.database.org/>. Accessed December 2009). Species previously reported from Guinea (Angel 1944. Bull. Mus. Nat. Hist. nat. Paris. 16:293–294), Ivory Coast (Barbault 1974. Bull. Soc. Zool. France. 99:345–361), and Gambia (Emms et al. 2007. Herpetol. Bull. 99:3–18).

Submitted by **SEBASTIEN TRAPE**, Laboratoire ECOLAG, Université Montpellier II, 34095 Montpellier Cedex 5, France; e-mail: sebastien_trape@yahoo.fr.

PLESTIODON FASCIATUS (Five-lined Skink). CANADA: ONTARIO: LAMBTON Co.: Village of Point Edward, ca. 60 m from shore of St. Clair River (42.992970°N, 82.410727°W; WGS 84). 20 May 2009. Daniel W. A. Noble and Jonathan D. Choquette. Verified by Ross D. MacCulloch. Royal Ontario Museum photographic vouchers (ROMdm 00251–00254). First municipality record and first observations in vicinity since 1932. Nearest record is historical from ca. 6 km S in the city of Sarnia, Ontario in 1932 (NMC 1740; Patch 1934. Copeia 1934:50–51). The Sarnia population (NHIC EO 16753) is considered extirpated by Ontario Ministry of Natural Resources Natural History Information Centre (NHIC). Records exist for only two other distant locations in Lambton County (both extant): Walpole Island (NHIC EO 1218) 45 km S and Pinery Provincial Park (NHIC EO 1212) 55 km NE. Three skinks (1 male,

1 female, 1 juvenile) were observed in a pile of logs, railway ties, and metal pipes in an isolated marina waste area. Two juveniles were observed 26 August and five juveniles on 11 September 2009 at same site. All were actively foraging or basking.

Submitted by **JONATHAN D. CHOQUETTE**, School of Environmental Design and Rural Development, University of Guelph, Guelph, Ontario N1G 2W1, Canada (e-mail: jchoquet@uoguelph.ca); **STEPHEN J. HECNAR**, Department of Biology, Lakehead University, Thunder Bay, Ontario P7B 5E1, Canada (e-mail: shecnar@lakeheadu.ca); **DANIEL W. A. NOBLE** (e-mail: nobled@uoguelph.ca), and **RONALD J. BROOKS**, Department of Integrative Biology, University of Guelph, Ontario N1G 2W1; (e-mail: rjbrooks@uoguelph.ca).

SQUAMATA — SNAKES

CROTALUS HORRIDUS (Timber Rattlesnake). USA: FLORIDA: VOLUSIA Co.: Lake Woodruff National Wildlife Refuge (29.105°N, 81.36697°W; WGS84). 20 July 2009. Wesley Allie and Wyatt Brouillard. Florida Museum of Natural History (UF 155622). Verified by Kenneth Krysko. First state record. Extends distribution ca. 61.2 km SE of nearest locality, Silver River State Park (Ashton and Ashton 1988. Handbook of Reptiles and Amphibians of Florida. Part One: The Snakes. Revised edition. Windward Publ. Co., Miami, Florida. 176 pp.).

Submitted by **W. BOYD BLIHOVDE** (e-mail: boyd_blihovde@fws.gov) and **KELLY A. BRADY**, Lake Woodruff National Wildlife Refuge, 2045 Mud Lake Road, DeLeon Springs, Florida 32130, USA (e-mail: kbrady@stetson.edu).

EUNECTES MURINUS (Green Anaconda). BRAZIL: PIAUÍ: Municipality of Ilha Grande, Delta of Parnaíba River, locality of Saquim (2.775656°S, 41.804756°W; datum WGS84). 13 October 2008. E. B. de Andrade and P. da C. Silva. Coleção Herpetológica do Museu Paraense Emílio Goeldi, Belém, Brazil (MPEG 23.36). Verified by T. C. Ávila-Pires. Species distributed in Trinidad Island, Venezuela, Colombia, Peru, Ecuador, Guiana, French Guiana, Suriname, and Brazil (Amazon Basin, central, and northeastern Brazil (Cunha and Nascimento 1993. Bol. Mus. Para. Emílio Goeldi, sér. Zool. 9[1]:1–191; Freitas and Silva 2007. A Herpetofauna das Caatingas e Áreas de Altitudes do Nordeste Brasileiro. 384 pp.). First record for state of Piauí, extending range ca. 332 km NE from the municipality of Arari, state of Maranhão (Cunha and Nascimento 1993, *op. cit.*) and ca. 390 km W from the municipality of Aquiraz, state of Ceará (Mendonça et al. 2009. Herpetol. Rev. 40:238).

Submitted by **ROBERTA R. DA SILVA-LEITE**, Universidade Federal do Piauí, Campus Ministro Reis Velloso, Av. São Sebastião nº 2819, Parnaíba, Piauí, Brazil, CEP 64202-020 (e-mail: Roberta.ufpi@gmail.com); **IGOR JOVENTINO ROBERTO**, Aquasis - Associação de Pesquisa e Preservação de Ecossistemas Aquáticos, Programa Biodiversidade. Praia de Iparana s/n, SESC Iparana, Caucaia, Ceará, Brazil, CEP 61627-010; **DANIEL LOEBMANN**, Departamento de Zoologia, Instituto de Biociências, Universidade Estadual Paulista, Rio Claro, São Paulo, Brazil, Caixa Postal 199, CEP 13506-970; **THIAGO DA SILVA NASCIMENTO**, Universidade Federal do Piauí, Campus Ministro Reis Velloso, Av. São Sebastião nº 2819, Parnaíba, Piauí, Brazil, CEP 64202-020; **PEDRO DA COSTA**