

JOSEPHI NICOLAI
LAURENTI
AUSTRIACI VIENNENSIS
SPECIMEN
MEDICUM,
EXHIBENS
SYNOPSISIN REPTILIIUM
EMENDATAM
CUM EXPERIMENTIS CIRCA VENENA
ET ANTIDOTA REPTILIIUM AUSTRIACORUM.



VIENNÆ,
TYP. JOAN. THOM. NOB. DE TRATTNERN,
CÆS. REG. AULÆ TYPOGR. ET BIBLIOP.

1768.

CVIII. *Seps ruber. Tab. III. fig. 3.*

DIAGN.

Collario soluto; scutis intermediis subrotundis; lateribus fuscis, dorso ruffo distinctissimo.

CIX. *Seps cærulescens. Tab. I. fig. 3.*

DIAGN.

Collario soluto; scutis intermediis subrotundis; corpus cæruleo - margaritaceum; latera ocellata triplici utrinque serie.

CX. *Seps varius. Tab. III. fig. 2.*

DIAGN.

Collario soluto, ruffo; scutis intermediis subrotundis; corpore viridi, punctis brunis variegato.

CXI. *Seps viridis. Seba Tom. II. Tab. CIII. fig. 4.*

DIAGN.

Collario soluto cæruleo; scutis intermediis subrotundis; corpore elegantissime viridi.

Habitat in aggeribus lapidosis sabulosis; in pratis sterilibus apricis; in muris & mœnibus.

Var. β. Ex viridi cærulescens.

γ. Nigris maculis variegata.

Hospitatur utraque in Museo Turriano.

Ordines scutorum abdominalium numero indefinito.

CXII.